

# Analysis of Bromine-Mercury Reactions in Flue Gas

Brydger Cauch, Paula A. Buitrago,  
Geoffrey D. Silcox, JoAnn S. Lighty  
Chemical Engineering, University of Utah

Connie L. Senior  
Reaction Engineering International

Andrew O'Palko  
U. S. Department of Energy  
National Energy Technology Laboratory  
DOE Award Number DE-FG26-06NT42713

2008 June 11


# Outline

- Objectives
- Introduction - Thermodynamics
- Kinetics and modeling of Hg-Br reactions
- Experimental
  - Homogeneous oxidation by Cl and Br
  - Effects of T profile, NO, Br, reactor surface area
  - Fixed-bed reactor
- Conclusions

# Project Objectives

- Develop validated models that allow prediction of extent of reaction between halogens and mercury in coal combustion applications.
- Include homogeneous and heterogeneous reactions.

# Introduction – Mercury Deposition



Ice core data on total deposited mercury (Wind River Mountains, Wyoming).

<http://toxics.usgs.gov/pubs/FS-051-02/>




# Introduction - Coal Composition


Sample Description	Elkhor/Hazard low S bit	Pittsburgh med S bit	Illinois 6 high S bit	Wyodak PRB	Wyodak PRB	Ohio 5,6,7 high S bit	ND Lignite lignite
<b>ANALYSIS (As Received):</b>							
Carbon	74.87	<b>76.62</b>	67.70	53.20	<b>51.19</b>	71.07	38.57
Hydrogen	4.59	<b>4.80</b>	4.73	4.59	<b>3.64</b>	4.81	2.60
Oxygen	8.38	<b>6.91</b>	9.19	20.74	<b>12.29</b>	8.10	12.52
Nitrogen	1.43	<b>1.48</b>	1.18	0.83	<b>0.72</b>	1.37	0.42
Sulfur	0.82	<b>1.64</b>	3.60	0.22	<b>0.32</b>	2.62	0.63
Ash	7.41	<b>7.01</b>	10.26	7.36	<b>6.03</b>	9.70	9.38
Moisture	2.33	<b>1.44</b>	3.31	13.06	<b>25.81</b>	2.33	35.88
Total	99.83	<b>99.89</b>	99.96	100.00	<b>100.00</b>	100.00	100.00
Hg, ug/g	0.13	<b>0.11</b>	0.22	0.19	<b>0.13</b>	0.15	0.13
Cl, ug/g	1660	<b>976</b>	338	**	<b>26</b>	974	36
Br, ug/g	25.0	<b>17.0</b>	3.7	2.4	<b>1.2</b>	23.0	1.9
Cl/Br	66	<b>57</b>	91	--	<b>22</b>	42	19

Analyses from DOE Toxics program. Equilibrium calculations use Pittsburgh bituminous, 3 % O<sub>2</sub>.

# Introduction – Gas-phase Thermodynamics of Cl and Br




HCl is dominant species at all T.




$\text{Br}_2$  is dominant species below 400°C.

# Introduction - Thermo of Hg/Cl, Hg/Br







$\text{HgCl}_2$  stable below 550°C.



$\text{HgBr}_2$  stable below 500°C.  
Chlorine excluded from this calculation.

# Kinetics - Hg/Br Reactions

- Theoretical and experimental studies support the following primary homogeneous oxidation pathways






# Kinetics - Hg/Br reactions

- Bromine-impregnated activated carbon shows 40-80% capture of mercury with hot-side ESP and >90% capture of mercury with cold-side ESP in low-halogen flue gas

# Kinetics and Modeling

- Modeling of homogeneous Hg-Cl and Hg-Br reactions
  - 468 reactions, 127 species
  - NO<sub>x</sub>, SO<sub>x</sub>, Br, Cl chemistry
  - Br chemistry from NIST website
  - Br-Hg chemistry developed for this work
  - 3% excess air, Pittsburgh bituminous

# Kinetics and Modeling




Typical time-  
temperature  
history in boiler



# Kinetics and Modeling

- Predicted homogeneous oxidation by halogens is negligible




- Equivalent coal halogens:
  - 70 ug/g Cl
  - 1, 60, 25 ug/g Br
- Equivalent flue gas halogens (as HX):
  - 5.7 ppmv HCl
  - 0.38, 2.3, 4.6 ppmv HBr
- Cl/Br molar ratios:  
150, 2.5, 1.2




# Quartz Tubular Reactor

- 300 W (1000 Btu/h), methane-fired




# Quartz Tubular Reactor

- Temperature profiles



# Sample Conditioning System




# Sample Conditioning System

- Sodium thiosulfate added to KCl impinger to prevent oxidation of  $\text{Hg}^0$  by hypochlorous acid.
  - $\text{Cl}_2 + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{HCl}$
  - $\text{HOCl} + \text{HCl} + \text{Hg}^0 \rightarrow \text{HgCl}_2 + \text{H}_2\text{O}$
  - $\text{Na}_2\text{S}_2\text{O}_3 + 4 \text{Cl}_2 + 5 \text{H}_2\text{O} \rightarrow 2\text{NaHSO}_4 + 8 \text{HCl}$
  - $\text{Na}_2\text{S}_2\text{O}_3 + 2 \text{HCl} \rightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{S} + \text{SO}_2$




# Oxidation by chlorine




# Oxidation by Bromine – Effect of Concentration

HQ (450 K/s), 1.5% O<sub>2</sub> (dry), 30 ppm NO (dry)




# Oxidation by Bromine – Effect of T Profile

LQ (220 K/s), 1.5% O<sub>2</sub> (dry), 30 ppm NO (dry)




# Oxidation by Bromine – Effect of NO

HQ (450 K/s), 1.5% O<sub>2</sub> (dry), 500 ppm NO (dry)





# Oxidation by Bromine – Effect of Reactor Surface Area

HQ (450 K/s), 1.5% O<sub>2</sub> (dry), 30 ppm NO (dry), Tube Bundle



# Future Work

- Heterogeneous oxidation on carbon
- Modeling – homogeneous and heterogeneous reactions



# Conclusions

- From thermodynamics,  $\text{Br}_2$  and  $\text{HCl}$  are dominant species at  $300^\circ\text{C}$ .
- $\text{HgCl}_2$  and  $\text{HgBr}_2$  are stable below about  $500^\circ\text{C}$ .
- Currently available kinetic models for homogeneous oxidation do not agree with our data.
- Sodium thiosulphate is essential for sample conditioning in absence of  $\text{SO}_2$ .

# Conclusions

- Homogeneous oxidation by chlorine is about 5% at 300 ppm Cl as HCl.
- Homogeneous oxidation by bromine is about 50% at 20 ppm Br as HBr.
- Homogeneous oxidation by chlorine is insensitive to quench rate while oxidation by bromine increases with lower quench rate.



# Conclusions

- Homogeneous oxidation of mercury by halogens is unaffected by NO and SO<sub>2</sub>.
- Extents of oxidation are insensitive to interior, quartz reactor surface area which implies that we are measuring homogeneous oxidation.